

Multimode electron-phonon coupling in the two-dimensional $t - J$ -Holstein model"

D. Poilblanc^{1,2}, T. Sakai¹, D. J. Scalapino² and W. Hanke^{3,2}

¹*Laboratoire de Physique Quantique, Université Paul Sabatier, 31062 Toulouse, France*

²*Department of Physics, University of California Santa Barbara, CA 93106*

³*Institute for Physics, University of Wurzburg, D-8700 Wurzburg, Germany*

(October 95)

Polaronic effects are investigated in the two dimensional t - J model (describing Cu spins and holes of the planes in high- T_c cuprates) in the presence of in-plane dynamic oxygen breathing modes. We focus on the effect of dynamic multi-phonon modes. Spin and charge structure factors are computed. We found evidence for self-localisation of the holes and *increased* short range antiferromagnetic spin-spin correlations between the remaining spins. This effect persists when all the in-plane oxygen vibration degrees of freedom are taken into account. This suggests that dynamic phonons could play a key role in stabilizing inhomogeneous stripe phases or domain wall structures.

PACS Numbers: 71.27.+a, 71.38.+i, 74.20.Mn, 74.25.Kc

The interplay of the electron-phonon interaction and localization, charge density wave (CDW) and spin density wave (SDW) correlations in strongly interacting electron systems is an interesting issue. In the static Born-Oppenheimer $m \rightarrow \infty$ limit, Zhong and Schüttler [1] studied Hubbard and t - J models with an onsite Holstein electron-phonon apical oxygen coupling. They found on Hubbard ($\sqrt{10} \times \sqrt{10}$) and t - J (4×4) clusters that in the nearly half-filled case, in the presence of a strong onsite Coulomb repulsion and short range antiferromagnetic spin correlations, there were enhanced polaronic self-localization effects. In a recent paper A. Dobry et al. [2] investigated the effects of a single dynamic $\mathbf{q}_0 = (\pi, \pi)$ phonon mode coupled to a strongly correlated two-dimensionnal (2D) electron system modeled by a t - J Hamiltonian. For cuprates superconductors, this (π, π) mode corresponds to a planar breathing phonon involving correlated displacements of the in-plane oxygens. For $\sqrt{10} \times \sqrt{10}$ clusters at one-quarter filling A. Dobry et al. [2] found evidence for a Charge Density Wave (CDW) instability at moderate values of the electron-phonon coupling. In addition, lattice deformations and CDW instabilities have been found in the one dimensional Holstein [3] and t - J -Holstein [4] models.

While both the Born-Oppenheimer and the dynamic single-mode approximations provide insight into this problem, one would like to examine the dynamic multi-phonon mode system. In particular, although the truncation of the Hilbert space of the phonon mode is well controlled, the single mode approach of Ref. [2] nevertheless assumes that in-plane oxygens oscillate in phase. However, there is experimental evidence that these modes have a small dispersion in the Brillouin zone, both in $La_{2-x}Sr_xCuO_4$ or in bilayer $YBa_2Cu_3O_{6+x}$ materials [5,6]. In other words, the oxygen vibrational modes are very localized and can be considered as independent. In the following, we investigate the influence of taking into account a *finite* number M of oxygen modes by perform-

ing exact diagonalisation studies of finite clusters. First, a simple model involving the coupling of a t - J Hamiltonian with two independent $\mathbf{q}_0$ breathing modes related to different sets of in-plane oxygen atoms is considered. Various signatures of a tendency towards hole localization (which could lead to CDW or domain wall structures) and the enhancement of short-range antiferromagnetic correlations between the remaining spins are found. We reach similar conclusions on a 2×2 lattice when all the in-plane oxygen vibration degrees of freedom are taken into account.

We start with a generalisation of the t - J -Holstein Hamiltonian [7],

$$H = -t \sum_{\langle \mathbf{i}, \mathbf{j} \rangle, \sigma} (\tilde{c}_{\mathbf{j}, \sigma}^\dagger \tilde{c}_{\mathbf{i}, \sigma} + \tilde{c}_{\mathbf{i}, \sigma}^\dagger \tilde{c}_{\mathbf{j}, \sigma}) + J \sum_{\langle \mathbf{i}, \mathbf{j} \rangle} (\mathbf{S}_{\mathbf{i}} \cdot \mathbf{S}_{\mathbf{j}} - \frac{1}{4} n_{\mathbf{i}} n_{\mathbf{j}}) + \Omega \sum_{\mathbf{i}, \delta} (\frac{p_{\mathbf{i}, \delta}^2}{2m} + \frac{1}{2} m \Omega^2 x_{\mathbf{i}, \delta}^2) + g \sum_{\mathbf{i}, \delta} x_{\mathbf{i}, \delta} (n_{\mathbf{i}}^h - n_{\mathbf{i}+\delta}^h) \quad (1)$$

where $\tilde{c}_{\mathbf{i}, \sigma}^\dagger$ is the usual hole creation operator, $n_{\mathbf{i}}$ and $n_{\mathbf{i}}^h$ are the electron and hole local densities respectively, m is the oxygen ion mass, Ω is the phonon frequency and $\delta = \mathbf{x}, \mathbf{y}$ differentiates the bonds along the x- and y-direction respectively. Throughout energies are measured in unit of the hopping integral t . Note that the electron-phonon g-term involves the coupling of each copper hole with the displacements of the four neighboring oxygens $x_{\mathbf{i}, \delta}$ and $x_{\mathbf{i}-\delta, \delta}$. For the purpose of our discussion it is convenient to re-write the electron-phonon coupling in momentum space,

$$H_{e-ph} = \Omega \sum_{\mathbf{q}, \delta} (b_{\mathbf{q}, \delta}^\dagger b_{\mathbf{q}, \delta} + \frac{1}{2}) + \frac{1}{\sqrt{2N}} \sum_{\mathbf{q}, \delta} \sum_{\mathbf{k}, \sigma} \lambda_{\mathbf{q}, \delta} (b_{\mathbf{q}, \delta}^\dagger + b_{-\mathbf{q}, \delta}) \tilde{c}_{\mathbf{k}-\mathbf{q}, \sigma}^\dagger \tilde{c}_{\mathbf{k}, \sigma} \quad (2)$$

where N is the number of sites, $b_{\mathbf{q},\delta}^\dagger$ are the usual boson creation operators, $\lambda_{\mathbf{q},\delta} = \lambda_0 \sin \frac{\mathbf{q} \cdot \delta}{2}$ and $\lambda_0 = g \sqrt{\frac{4}{m\Omega}}$. The Hamiltonian of Ref. [2] is obtained from Eq. (2) by (i) keeping only a single δ , $\mathbf{q} = \mathbf{q}_0$ mode and (ii) replacing the sum $\frac{1}{\sqrt{2N}} \sum_{\mathbf{q},\delta}$ by unity. This is *a priori* justified by the fact that the coupling constant $\lambda_{\mathbf{q},\delta}$ is maximum at wave vector $\mathbf{q}_0$. However, assuming in-phase oxygen displacements might overestimate cooperative effects of the phonons. On the other hand, a complete treatment of hamiltonian (2) requires to take the $M=2N$ oxygen phonons into account.

The first step to go beyond the single mode approximation (SMA) of Hamiltonian (2) is to consider two independent bosons $b_{\mathbf{q}_0,\mathbf{x}}$ and $b_{\mathbf{q}_0,\mathbf{y}}$ ($M=2$) for the horizontal ($\delta = \mathbf{x}$) and vertical bonds ($\delta = \mathbf{y}$). In this case, the sum $\frac{1}{\sqrt{2N}} \sum_{\mathbf{q},\delta}$ is replaced by $\frac{1}{2} \sum_{\delta}$ and only two $\mathbf{q} = \mathbf{q}_0$ modes are retained. The ground state (GS) energies $E_0(\lambda_0)$ of the single and double mode approximations (DMA) of Hamiltonian (2) are calculated on a $N = \sqrt{20} \times \sqrt{20}$ cluster doped with two holes by truncating the (strictly speaking infinite) Hilbert space of the phonon mode(s) to a finite number of bosonic levels N_{ph} . This can be achieved, for example, by keeping only a finite number of levels $n_{ph} + 1$ for each oscillator. In other words, a maximum number of n_{ph} local phonons could be excited for each oxygen mode. Then the total number of degrees of freedom for the bosons becomes $N_{ph} = (n_{ph} + 1)^M$ where M is the number of modes retained ($M=1$ for SMA, $M=2$ for DMA and $M=2N$ for the full problem).

The GS energies $E_0(\lambda_0) - E_0(0)$ for $M=1$ and $M=2$ are displayed in Figs. 1(a,b) in a regime ($J = 0.3$) where, in the absence of phonons, spin fluctuations leads to a small effective attraction between the holes [8]. The dependence on the number of levels $n_{ph} + 1$ is shown in Fig. 1(a). For $M=1$, as in the quarter filled case considered in Ref. [2], the truncating procedure is well controlled and the GS energy decreases with N_{ph} and typically converges for $N_{ph} \sim 20$ or 30. When more than a single mode is included, it is clear from Fig. 1(a) that n_{ph} (and not N_{ph}) is the key parameter controlling the convergence of the GS energy. We then expect that a similar number of levels per phonon mode (~ 15) is still required to fully account for the vibrational degrees of freedom when $M > 1$. Hence, disk space limitations clearly restrict the study of the full hamiltonian (2) (with $M = 2N$) to small 2×2 clusters (see below). However, useful informations can still be obtained with $n_{ph} \sim 2-5$ since most physical properties are smooth monotonic functions of n_{ph} . Fig. 1(a) suggests that polaronic effects are stronger when the number of independent modes (M) is increased. As seen from Fig. 1(b), for increasing λ_0 both models show a crossover from an adiabatic regime (the energy depends weakly on the coupling strength) to a polaronic regime. We, indeed, expect that, for a sufficiently large value of the electron-phonon coupling, the

system will start to form large hole polarons. By comparing the set of data for the two models we observe that the onset λ_{co} corresponding to polaron formation might not depend too strongly on the truncation parameter M (or n_{ph}) although for $\lambda_0 > \lambda_{co}$ polaronic effects increase with M .

In addition to the formation of polarons, the electron-phonon coupling (treated in the SMA) also favors the configurations where the holes (polarons) stay dynamically on the same sublattice. Such a polaron-polaron interaction can eventually lead to a CDW instability as, for example, it is found at quarter filling [2] in the SMA or to domain walls formation. To complete our study, we address now the issue of the spin and charge correlations at low hole doping (10%) and study the influence of the number of phonon modes on the (π, π) structure factors. We shall consider here only the case of commensurate (π, π) charge fluctuations since the density is too close to half filling and the system probably not large enough to accomodate any incommensurate order. The charge and spin structure factors at momentum (π, π) are given by

$$S_c(\mathbf{q}_0) = \langle \Psi_0 | \left(\sum_{\mathbf{i}} (-1)^{(i_x + i_y)} n_{\mathbf{i}}^h \right)^2 | \Psi_0 \rangle \quad (3)$$

$$S_s(\mathbf{q}_0) = \langle \Psi_0 | \left(\sum_{\mathbf{i}} (-1)^{(i_x + i_y)} S_{\mathbf{i}}^z \right)^2 | \Psi_0 \rangle \quad (4)$$

where $|\Psi_0\rangle$ is the GS wavefunction calculated both in the SMA and DMA cases. $S_c(\mathbf{q}_0)$ is shown in Fig. 2 (a) as a function of the electron-phonon coupling. For a single mode (SMA) a significant increase of the charge structure factor occurs around $\lambda_0 = 0.45$ indicating a tendency towards CDW formation or at least towards some localisation of the charge carriers. Interestingly enough, Fig. 2 (b) shows that this transition gives rise to a simultaneous increase of the spin-spin antiferromagnetic correlations. This phenomenon might be related to a sudden increase of the polaronic mass or to the localisation of the holes in some kind of (fluctuating) CDW state or stripe (domain wall) phase. Less mobile holes clearly lead to a weaker perturbation of the antiferromagnetic background, hence leading to enhanced short range spin-spin antiferromagnetic correlations. It should be noticed that, on the contrary, for quarter filling [2] the CDW transition is followed by a decrease of the spin correlations. When two oxygen vibrational modes are considered ($M=2$), Fig. 2 (a) and (b) show that the increases of the charge and spin structure factors (indicating some hole localization) occur for approximately the same couplings than for $M=1$.

We finish our study by the full treatment of all the in-plane oxygen degrees of freedom as seen in Fig. 3(a,b). Although the system size is smaller in this case (2×2) the results are qualitatively similar. Fig. 3(b) clearly shows a large increase of the spin structure factor at λ_{co} . This increase becomes sharper and sharper when more phononic levels are included (eventhough the GS energy is not fully converged, see Fig. 3(a)). Note that the crossover value seems to be smaller for $n=0.75$ ($\lambda_{co} \sim 0.3$)

than for $n=0.9$ ($\lambda_{co} \sim 0.4$).

The previous results can be physically interpreted as follows; In the SMA the oxygen atoms are constrained to oscillate in phase leading to a dynamical potential which favors the occupation by the carriers of one of the two sublattices. As discussed in Ref. [2] this can lead to a $\mathbf{q}_0$ CDW instability as can be seen for example in a mean-field treatment. The DMA is the simplest way to introduce some dephasing between the harmonic oscillators. In the extreme case of out-of-phase motions of the two oxygen sets the effective potential for the holes exactly vanishes. Therefore, one would naively expect that the DMA or the complete treatment of hamiltonian (2) lead to a weaker tendency towards CDW than the SMA. On the contrary, our results show that, in these more reliable treatments, the electron-phonon coupling still leads to strong self-localization effects [9]. More importantly, they also suggest that dynamic in-plane phonons could help to stabilize inhomogeneous phases [10,11] where holes are squeezed into stripes or clumps and where the remaining antiferromagnetic regions have enhanced $S_s(\pi, \pi)$ correlations.

D. P. thanks G. Zawatsky for insightful discussions on experimental aspects and J. Riera for interesting correspondence. *Laboratoire de Physique Quantique, Toulouse* is *Unité de Recherche Associé au CNRS No 505*. DP and DJS acknowledge support from the EEC Human Capital and Mobility program under Grant No. CHRX-CT93-0332 and the National Science Foundation under Grant No. DMR92-25027 respectively. We also thank IDRIS, Orsay (France), for allocation of CPU time on the C94 and C98 CRAY supercomputers.

Wellein, H. Röder and H. Fehske, preprint.

- [10] V. J. Emery, S. A. Kivelson and H. Q. Lin, Phys. Rev. Lett. **64**, 475 (1990).
- [11] D. Poilblanc and T. M. Rice, Phys. Rev. B **39** 9749 (1989).

FIG. 1. Pair energy calculated on a $\sqrt{20} \times \sqrt{20}$ t-J cluster with 2 holes for $J = 0.3$ and $\Omega = 0.2$. Open (solid) symbols correspond to a single (double) (π, π) breathing mode. (a) Energy vs number of phononic levels included in the calculations. (b) Energy vs coupling strength λ_0 . The number of phononic levels considered is indicated on the plot for each case (single and double modes).

FIG. 2. Charge (a) and spin (b) structure factors at momentum (π, π) vs electron-phonon coupling calculated on a $\sqrt{20} \times \sqrt{20}$ t-J cluster doped with 2 holes for $J = 0.3$ and $\Omega = 0.2$. Open (solid) symbols correspond to a single (double) (π, π) breathing mode. $N_{ph} = 25$ phononic levels were kept in the calculation.

FIG. 3. GS energy vs n_{ph} (a) and spin structure factor at momentum (π, π) vs electron-phonon coupling (b) calculated on a 2×2 t-J cluster doped with 1 hole for $J = 0.3$ and $\Omega = 0.2$. The total 8 independant oxygen modes have been considered.

-
- [1] J. Zhong and H.-B. Schüttler, Phys. Rev. Lett. **69**, 1600 (1992).
 - [2] A. Dobry, A. Greco, S. Koval and J. Riera, Phys. Rev. B, in press, Nov. 1995.
 - [3] J. E. Hirsch and E. Fradkin, Phys. Rev. B **27**, 4302 (1983).
 - [4] R. Fehrenbacher, Phys. Rev. B **49**, 12230 (1994).
 - [5] W.H. Weber et al., Phys. Rev. B **38**, 917 (1988); L. Pintschovius et al., Physica C **185-189**, 156 (1991).
 - [6] L. Pintschovius et al., in Physical Properties of High Temperature Superconductors IV, ed. D.M. Ginsberg, World Scientific (1994).
 - [7] For derivations of low energy effective electron-phonon Hamiltonians from 3-band Hubbard models see e.g. K. von Szczepanski and K. Becker, Z. Phys. B **89**, 327 (1992).
 - [8] Y. Hasegawa and D. Poilblanc, Phys. Rev. B **40** 9035 (1989); *ibid* **48**, 3368 (1993); D. Poilblanc, J. Riera and E. Dagotto, Phys. Rev. B **49**, 12318 (1994).
 - [9] Large polaronic effects have also been found recently in the case of apical oxygen out-of plane motions; G.





